

ON THE PRE-IMAGE OF A POINT UNDER AN ISOGENY

JONATHAN REYNOLDS

October 1, 2008

Abstract

Given a rational point on a curve in a rational isogeny class, a natural question concerns the field of definition of its pre-images. The multiplication by m endomorphism is a powerful and much-used tool. The pre-images for this map are found by factorizing a monic polynomial of degree m^2 . For $m = 2$, Everest and King gave examples where the existence of a quadratic factor coincided with the existence of a rational pre-image via a 2-isogeny. Nelson Stephens asked if this always happens and the question is answered in the affirmative. It is also shown that the analogue for $m = 3$ can only be false when there exists a rational point of order three and a small number of counterexamples are found. The results are proven over any field with characteristic not two or three.

1 Introduction

Given an elliptic curve $E/\mathbb{Q}$, the set of all curves E' isogenous to E over $\mathbb{Q}$ is finite (up to isomorphism) and is known as an isogeny class. Vélú's formulae [10] and the Weierstrass parameterization of the elliptic curve can be used to find an isogeny class. This is best illustrated in an algorithm developed by Cremona [2]. He has used his algorithm to produce tables of isogeny classes [3]. For each curve in the class, non-torsion generators of the Mordell-Weil group are also given. Methods which find isogeny classes over an arbitrary field have also been developed [4]. Given a curve in an isogeny class and a point in the Mordell-Weil group, the focus here is on where the pre-images of the point are defined.

Definition 1.1. *Let K be a field, E/K an elliptic curve, $P \in E(K)$ and $\phi : E' \rightarrow E$ an isogeny. Suppose that E' , ϕ and a point in $\phi^{-1}(P)$ are all defined over a finite extension L/K with $[L : K] < \deg \phi$. Then P is called magnified. If $L = K$ then P is called K -magnified.*

Suppose that $E/\mathbb{Q}$ is an elliptic curve. In [5] Everest, Miller and Stephens showed that if a rational non-torsion point is $\mathbb{Q}$ -magnified by $\phi : E' \rightarrow E$ then the corresponding elliptic divisibility sequence has only finitely many prime

power terms. In [7] it was shown that these finitely many terms correspond to S -integral points on E' , where S is given explicitly. By the main result in [6], the $\mathbb{Q}$ -magnified condition can be weakened to magnified when the isogeny is multiplication by two or three. Stephens asked if a non-torsion point is $\mathbb{Q}$ -magnified whenever it is magnified by doubling a point. The following theorem resolves this question.

Theorem 1.2. *Let E be an elliptic curve defined over a field K with $\text{char } K \neq 2$. If a non-torsion point $P \in E(K)$ is magnified by doubling a point then it is K -magnified.*

It should be noted that factorizing a monic quartic polynomial determines whether a point is magnified by doubling a point (see Section 2). This polynomial depends only on the point and the coefficients of a Weierstrass equation for E . For multiplication by three the conclusion has to be weakened.

Theorem 1.3. *Let E be an elliptic curve defined over a field K with $\text{char } K \neq 2, 3$. If a non-torsion point $P \in E(K)$ is magnified by tripling a point then either it is K -magnified or E has a non-trivial K -rational 3-torsion point.*

Listed in Table 1 are curves having a non-torsion rational point which is magnified by tripling a point but not $\mathbb{Q}$ -magnified. They were found using PARI/GP [9] and in Section 3 it is explained that such examples are rare.

2 Proofs of Theorems 1.2 and 1.3

Let K be a field with $\text{char } K \neq 2$. Let E/K be an elliptic curve with Weierstrass coordinate functions x and y . Let $P \in E(K)$ be a non-torsion point. The following observation plays an important role.

Lemma 2.1. *Suppose that E'/K is an elliptic curve and $\phi : E' \rightarrow E$ is an isogeny defined over K with $\phi(R) = P$. Then $K(x(R), y(R)) = K(x(R))$.*

Proof. Put $L = K(x(R))$ and $L' = K(x(R), y(R))$. Then $[L' : L] \leq 2$. Suppose that $[L' : L] = 2$ and choose $\sigma \in \text{Gal}(L'/L)$ to be non-trivial. Then $T = \sigma(R) - R$ is in the kernel of ϕ since $\sigma(\phi(R)) - \phi(R) = \mathcal{O}$. But $x(R + T) = x(R)$ so $R + T = \pm R$. Since P is non-torsion it follows that $\sigma(R) = R$ and $L' = L$. $\square$

Let $\psi_m, \theta_m \in K[E]$ be the standard division polynomials (see p. 39 of [1]). Define $\delta_m^P \in K[x]$ by $\delta_m^P = \theta_m - x(P)\psi_m^2$. Then δ_m^P is monic and has degree m^2 . The zeros of δ_m^P determine the values of $x(R)$ for which $mR = P$.

Proof of Theorem 1.2. Assume that P is magnified by doubling $R \in E(L)$. Using Lemma 2.1, $L = K(x(R))$. We may choose R so that $[L : K] \leq 2$. Suppose that $[L : K] = 2$ and choose $\sigma \in \text{Gal}(L/K)$ to be non-trivial. Then $T = \sigma(R) - R$ is a 2-torsion point since $\sigma(2R) - 2R = \mathcal{O}$. Also $T \in E(K)$ since $\sigma(T) = -T$. Using this torsion point, we can construct an elliptic curve E'/K and a 2-isogeny $\phi : E \rightarrow E'$ with $\ker \phi = \{\mathcal{O}, T\}$ (see p. 95 of [8]). Moreover, both ϕ

and its dual $\hat{\phi} : E' \rightarrow E$ are defined over K . Put $\phi(R) = Q$. It follows that $\sigma(Q) = \phi(\sigma(R)) = \phi(R + T) = \phi(R)$. Hence $Q \in E'(K)$ and $\hat{\phi}(Q) = P$. $\square$

Proof of Theorem 1.3. We may assume that δ_3^P has an irreducible factor f of degree 3. Let r_1, r_2, r_3 be the zeros of f and let R_1, R_2, R_3 be the corresponding solutions to $3R = P$. Then the splitting field L of f is $K(r_1, \sqrt{D})$, where D is the discriminant of f . Moreover, $\text{Gal}(L/K)$ is isomorphic to A_3 or S_3 . Let $\alpha = (1, 2, 3)$ be the generator of $\text{Gal}(L/K(\sqrt{D}))$ and put $T = \alpha(R_1) - R_1$. Suppose that E does not have a non-trivial K -rational 3-torsion point. Any permutation preserves $R_1 + R_2 + R_3 = P$ so it must equal $\mathcal{O}$. Thus $\alpha(T) = R_3 - R_2 = R_2 - R_1 = T$ and D is not a square in K . Choose $\sigma \in \text{Gal}(K(\sqrt{D})/K)$ to be non-trivial. Since $\sigma(T) + T$ is a K -rational 3-torsion point, $x(T) \in K$. Using Vélú's formulae [10], we can construct an elliptic curve E'/K and a 3-isogeny $\phi : E \rightarrow E'$ with kernel $\{\mathcal{O}, T, -T\}$. Moreover, both ϕ and its dual $\hat{\phi} : E' \rightarrow E$ are defined over K . Now $\phi(R_2) = \phi(R_1 + T)$ and $\phi(R_3) = \phi(R_1 - T)$. Hence $\phi(R_1)$ is fixed by $\text{Gal}(L/K)$. $\square$

3 Computations

Examples where δ_3^P factorizes (over K) but P is not K -magnified are relatively hard to find. Let $K = \mathbb{Q}$. The first 12 of Cremona's "generators" tables from [3] were considered. There are 22,962 pairs (E, P) such that δ_3^P factorizes (over $\mathbb{Q}$). In all but 14 of these pairs P is $\mathbb{Q}$ -magnified by a 3-isogeny, which can be constructed using Vélú's formulae [10] and a rational zero of ψ_3 .

Table 1: Curves having magnified points which are not $\mathbb{Q}$ -magnified

N	C	$\#$	N	C	$\#$	N	C	$\#$
17739	g	1	47526	f	1	70470	m	2
19926	l	2	49818	j	1	92055	u	1
26730	y	2	57222	bw	2	113866	d	1
39710	z	1	62814	r	1	119646	dd	2
45662	h	1	64395	f	1			

The 14 counterexamples found are listed in Table 1. They are given in the same format as Cremona uses, where N is the conductor, C is the isogeny class and $\#$ is the number of the curve in the class. Each curve listed has rank 1 and P is taken to be the generator which Cremona gives. Moreover, they each lie in an isogeny class of size two. Let Q be a generator for the other curve. Denote by $\hat{h}$ the canonical height (see [8]). If P is $\mathbb{Q}$ -magnified by a d -isogeny then it is the image of $mQ + T$, for some non-zero integer m and torsion point T . This implies $dm^2\hat{h}(Q) = \hat{h}(P)$. But in fact, $\hat{h}(Q) = 3\hat{h}(P)$. Thus P is not $\mathbb{Q}$ -magnified.

It is worth noting that for all of the pairs (E, P) considered and for all odd primes $l \leq 7$, if δ_l^P factorizes (over $\mathbb{Q}$) then ψ_l has a factor of degree at most $(l - 1)/2$.

References

- [1] I. Blake, G. Seroussi and N. Smart, *Elliptic Curves in Cryptography* (Cambridge University Press, 1999).
- [2] J. E. Cremona, *Algorithms for Modular Elliptic Curves* (Cambridge University Press, 1997).
- [3] J. E. Cremona, Elliptic Curve Data, updated 2006-09-24, <http://www.warwick.ac.uk/~masgaj/ftp/data/>.
- [4] J. E. Cremona and M. Watkins, Computing Isogenies of Elliptic Curves, preprint available from <http://www.maths.bris.ac.uk/~mamjw/>.
- [5] G. Everest, V. Miller and N. Stephens, Primes Generated by Elliptic Curves, *Proc. AMS*, **132** (2004) 955–963.
- [6] G. Everest and H. King, Prime Powers in Elliptic Divisibility Sequences, *Mathematics of Computation*, **74** (2005) 2061–2071.
- [7] G. Everest, J. Reynolds and S. Stevens, On the denominators of rational points on elliptic curves, *Bull. London Math. Soc.* **39**(5) (2007) 762–770.
- [8] D. Husemöller, *Elliptic Curves* (Springer, 2004).
- [9] The PARI Group, PARI/GP, version 2.3.2 (2005), <http://pari.math.u-bordeaux.fr/>.
- [10] J. Vlu, Isognies entre courbes elliptiques, *C. R. Acad. Sci. Paris*, **273** (1971) 238–241.